



Standardized and simple sub-fractionation of human plasma reveals enrichment of many low abundant hydrophobic proteins

Jessen, Flemming; Wulff, Tune

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Jessen, F., & Wulff, T. (2012). *Standardized and simple sub-fractionation of human plasma reveals enrichment of many low abundant hydrophobic proteins*. Poster session presented at 9th Siena Meeting From Genome to Proteome, Siena, Italy.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Standardized and simple sub-fractionation of human plasma reveals enrichment of many low abundant hydrophobic proteins

Flemming Jessen and Tune Wulff

National Food Institute, DTU, Kgs. Lyngby, Denmark (fjes@food.dtu.dk)

Aim

Sub-fractionation of blood plasma


Analysis

Human plasma (2 µl) and proteins extracted from 400 µl plasma were analysed by 2DE (pI: 4-7 and 3-10; T: 12 %; Colloidal Coomassie staining)

Background:


A main characteristic of plasma proteomics is the *10 order of magnitude problem*, causing that low abundance proteins are hard to visualise using 2D-gel electrophoresis (2DE). This problem calls for sub-fractionation to simplify the protein composition of the plasma sample, making it possible to detect lower abundant plasma proteins. Direction of strategies varies from depletion of high abundant proteins to enrichment of low abundant proteins using different approaches. However, a major concern when sub-fractionating is the reproducibility.

Extracted proteins




#	Protein ID (MALDI-TOF-TOF)
1	Actin, cytoplasmic 1
2	Actin, acrosomal process isoform
3	Adipocyte plasma membrane-associated protein
4	Alpha-1-antitrypsin
5	Apolipoprotein A-I
6	Apolipoprotein E
7	Ficolin-3
8	Lipopolysaccharide-binding protein
9	Prenylcysteine oxidase 1
○	Apolipoprotein M
○	Haptoglobin-related protein
○	Apolipoprotein L1
○	Serum paraoxonase/arylesterase 1
○	Serum amyloid A

pI: 3-10




Blood plasma



Positions of extracted spots where no or very low abundance spots are present in a 2D-gel of plasma

Technical and biological variation



Data (normalised spot volumes) from 2DE analyses of 10 protein extracts made from 10 subjects (Biol+Tech) and 10 protein extracts made from a plasma pool (Technical)